


Table 2: De Novo Sequence Assemblies

Figure Plot	Sample [*]	% GC	Reference Size (bp)	Coverage Depth (x)	K-mer Size	Contig N50 Length (bp)	Number of Contigs	Scaffold N50 Length (bp)	Number of Scaffolds	Assembly Length (bp)
A	<i>L. monocytogenes</i> (1)	37.98	2,944,528	68.02	75	1,620,614	7	2,932,577	4	2,938,359
B	<i>L. monocytogenes</i> (2)	37.98	2,944,528	53.62	75	691,775	11	2,928,935	5	2,933,419
C	<i>M. ruber</i> (1)	63.38	3,097,457	55.43	75	203,917	26	3,001,248	9	3,100,592
D	<i>M. ruber</i> (2)	63.38	3,097,457	47.96	69	145,226	35	2,998,617	14	3,105,938
E	<i>P. heparinus</i> (1)	42.05	5,167,383	36.07	51	297,483	57	5,144,140	16	5,160,459
F	<i>P. heparinus</i> (2)	42.05	5,167,383	27.48	55	289,124	65	5,150,646	14	5,158,523
G	<i>K. pneumoniae</i> (1)	57.15	5,694,894	32.28	59	142,641	141	4,455,493	40	5,575,733
H	<i>K. pneumoniae</i> (2)	57.15	5,694,894	29.37	51	159,999	144	3,651,106	46	5,579,858
I	<i>B. cereus</i> (1)	35.50	5,432,652	24.24	43	90,460	122	1,710,854	24	5,384,726
J	<i>B. cereus</i> (2)	35.50	5,432,652	27.36	51	100,734	106	1,149,845	20	5,398,264
K	<i>R. sphaeroides</i> (1)	68.79	4,602,977	39.17	61	191,856	88	4,129,505	14	4,503,303
L	<i>R. sphaeroides</i> (2)	68.79	4,602,977	45.22	69	210,411	78	3,186,951	17	4,513,006
M	<i>E. coli</i> DH10B (1)	50.78	4,686,137	48.38	57	234,160	65	4,422,317	26	4,545,800
N	<i>E. coli</i> DH10B (2)	50.78	4,686,137	35.31	57	178,404	76	1,591,696	25	4,537,054
O	<i>M. tuberculosis</i> (1)	65.61	4,419,977	47.19	73	92,267	97	4,354,886	23	4,362,803
P	<i>M. tuberculosis</i> (2)	65.61	4,419,977	38.99	53	88,733	117	2,527,876	27	4,359,302
Q	<i>E. coli</i> MG1655 (1)	50.79	4,639,675	35.60	61	179,369	69	4,590,510	22	4,598,448
R	<i>E. coli</i> MG1655 (2)	50.79	4,639,675	35.43	55	198,090	64	4,595,933	18	4,603,074

^{*} The replicate number for each strain is denoted in parentheses.

Reference sizes include plasmids. Coverage depth denotes the coverage after adapter trimming, excluding reads < 25 bp in length after trimming. The assembly data presented correspond to contigs > 500 bp in length except where "scaffold" is denoted. Reference genomes were obtained from the NCBI index of genomes⁷.

Figure 3: De Novo Assemblies Compared to Reference Genomes



These plots demonstrate the concordance between the bacterial genome assemblies and the respective reference genomes. Table 2 defines the strains associated with plots A–R. Reference genomes are shown on the X-axes and assemblies are shown on the Y-axes. A red line represents a consistent segment between the assembly and the reference. A blue line represents a consistent (inverted) segment. The vertical lines separate chromosomes and plasmids within the reference. The horizontal lines separate scaffolds in the assemblies. The misassemblies shown in plots A, J, and N could be refined with further analysis. These plots were generated using MUMmer⁸.

AAAGAATGATAACAGTAACACACTTCTGTTAACCTTAAGATTACTTGATCCACTGATTCAACGTACCGTAAACGAACGTATCAATTGAGACTAAATATTAACGTACCATTAAAGAGCTACCGTCTTCTGTTAACCTTAAGATTACTTGATCCACTGATTCA
AATCAACGTACCGTAACGAACGTATCAATTGAGATTACTTGATCCACTGATTCAACGTACCGTAAACGAACGTATCAATTGAGACTAAATATTAACGTACCATTAAAGAGCTACCGTCTTCTGTTAACCTTAAGATTACTTGATCCACTGATTCA
AACGACGAAAGAATGATAACAGTAACACACTTCTGTTAACCTTAAGATTACTTGATCCACTGATTCAACGTACCGTAAACGAACGTATCAATTGAGACTAAATATTAACGTACCATTAAAGAGCTACCGTCTTCTGTTAACCTTAAGATTACTTGATCCACTGATTCA
TTAACGTACCATTAAAGAGCTACCGTCAACAGTAACACACTTCTGTTAACCTTAAGATTACTTGATCCACTGATTCAACGTACCGTAAACGAACGTATCAATTGAGACTAAATATTAACGTACCATTAAAGAGCTACCGTCAACGACGAAAGAATGAT
AAAGAATGATAACAGTAACACACTTCTGTTAACCTTAAGATTACTTGATCCACTGATTCAACGTACCGTAAACGAACGTATCAATTGAGACTAAATATTAACGTACCATTAAAGAGCTACCGTCTTCTGTTAACCTTAAGATTACTTGATCCACTGATTCA
AAGATTACTTGATCCACTGATTCAACGTAAAGATTACTTGATCCACTGATTCAACGTACCGTAAACGAACGTATCAATTGAGACTAAATATTAACGTACCATTAAAGAGCTACCGTCAACGACGAAAGAATGATAAC
AACGTATCAATTGAGACTAAATATTAACGTACCATTAAAGAGCTGTTAACCTTAAGATTACTTGATCCACTGATTCAACGTACCGTAAACGAACGTATCAATTGAGACTAAATATTAACGTACCATTAAAGAGCTACCGTCAACGACGAAAGAATGATAAC

